

Early-season Performance of Diluted Concord Grape Juice and Commercial Lures at Attracting Spotted Wing Drosophila and Effects of Juice Fermentation on Trap Captures

Jaime C. Piñero and Heriberto Godoy-Hernandez

Stockbridge School of Agriculture, University of Massachusetts

Commercial synthetic food-based lures are available to monitor spotted wing drosophila (SWD). While effective, most commercial lures are based on fermentation materials and consequently they attract a comparatively high number of non-target insects. Captures of unwanted insects hinders trap performance and increases sorting time. For the past two years we have been evaluating the attractiveness of diluted Concord grape juice, a low-cost and readily available material, to male and female SWD. When diluted at a ratio of 1:3 (= 1 part of grape juice and three parts of water), diluted grape juice showed to be three times more attractive to males and females than one commercial lure under field conditions (see Piñero and Foley, 2018). In addition, grape juice diluted at the 1:3 ratio attracted significantly fewer (about three times less) non-targets than one commercial lure, highlighting a potential greater selectivity of diluted Concord grape juice.

Here, we assessed the efficacy of traps baited with diluted Concord grape juice at monitoring SWD early in the season when compared to four commercial lures. A secondary objective was to elucidate the effects of grape juice fermentation, as influenced by aging, on SWD captures in traps. For all experiments, we assessed the selectivity of the materials evaluated by relating captures of SWD to non-target insects.

Materials & Methods

Experiment 1: Early-season SWD monitoring

using grape juice and commercial lures. Here, we quantified the ability of 1-quart traps baited with diluted grape juice at the 1:3 dilution ratio (thereafter referred to as ‘DGJ’) to detect the first adult SWD of the growing season, compared with 1-quart traps baited with commercial lures. Five olfactory treatments were evaluated (1) DGJ, (2) Scentry® lure, (3) AlphaScents® lure, (4) Trécé broad spectrum PEEL-PAK multi-component lure, and (5) Trécé high-selectivity 3-component lure. DGJ-baited traps received 6 oz. of this material whereas traps baited with commercial lures received 6 oz. of unscented soapy water as a drowning solution. On May 4, five traps (one trap of each treatment) were deployed at each of four locations: Deerfield (on cherry trees), Belchertown (cherry), Whately (raspberry), and Amherst (elderberry). Traps were deployed on May 4 and they were removed on July 8, 2020. Once a week, all traps were serviced and all captured insects were brought back to the laboratory, for identification. At each trap inspection session, DGJ was replaced. The commercial lures remained in place until traps were removed.

Experiment 2. Effects of DGJ fermentation. This field test was conducted during June and July, 2019, in a cherry block at the UMass Cold Spring Orchard. The objective of this evaluation was to quantify the response of male and female SWD to volatiles emitted by DGJ (1:3 ratio) that had been aged for either, 7 or 14 days, compared to fresh DGJ. The four treatments evaluated were (1) fresh DGJ, (2) 7-day old DGJ, (3) 14-days old

DGJ, and (4) water as control. DGJ was prepared in the laboratory using 1-gallon containers with a gauze secured with rubber bands to allow for air exchange. GJD was prepared three times a week, and aged inside a chamber at 77°F and 65-75% relative humidity. The baits were prepared following a schedule that allowed for simultaneous evaluations of aged materials. Each treatment was replicated 5 times (= 5 cherry trees). Twice a week, traps were serviced, and insects were collected and transported to the lab, for identification. At each inspection session, clean traps were re-baited with newly-prepared materials.

Results

Experiment 1: Early-season comparison of diluted grape juice and commercial lures. As shown in Table 1, traps baited with diluted grape juice (DGJ) and with Trécé lures captured the first adult SWD of the season. This occurred on May 25, 2020. On June 1, DGJ was the only treatment that attracted SWD to traps. On June 10, all treatments except for AlphaScents attracted SWD. Results from the June 17 trapping date showed that the Scentry lure attracted the highest number of SWD. The Trécé broad spectrum lure ranked second on this date. DGJ attracted the highest number of SWD on June 30 and on July 8 (Table 1).

The selectivity of each treatment is presented in Table 1 (see section ‘Ratio (other drosophilids to SWD)’). DGJ was the most selective treatment (lower values

Table 1. Early-season captures (expressed as average number per trap) of spotted wing drosophila (SWD) and non-target insects (other insects belonging to the SWD family) in traps baited with diluted grape juice (DGJ) and four commercial lures at four Massachusetts locations. Boxes highlighted in orange indicate the date of first SWD captures in 2020.

Treatment	May 11	May 18	May 25	June 1	June 10	June 17	June 22	June 30	July 8
SWD									
DGJ	0	0	0.7	0.3	0.8	4.8	1.3	8.3	16.8
AlphaScents	0	0	0	0	0	4.3	2.8	2.3	2.8
Scentry	0	0	0	0	2.0	11.3	3	3.3	1.5
Trécé broad spectrum	0	0	0.7	0	0.5	8.3	1.3	3.0	1.5
Trécé selective	0	0	1.0	0	0.3	2.0	0.5	2.0	0.3
Other drosophilids									
DGJ	0	0.7	16.7	11	3.3	6.8	1.8	14.3	32.3
AlphaScents	14.3	39.7	4	6.5	18.8	58.3	23.3	8.0	13.0
Scentry	2	27	66	11.8	5	35.5	64.5	6.0	12.0
Trécé broad spectrum	55.7	88.3	76	14	54	106.8	10.5	10.3	22.0
Trécé selective	3	9.7	30.3	4	5.3	9.0	1.7	2.0	3.8
Ratio* (other drosophilids to SWD)									
DGJ	---	---	23.9	36.7	4.1	1.4	1.4	1.7	1.9
AlphaScents	---	---	---	---	---	13.6	8.3	3.5	4.6
Scentry	---	---	---	---	2.5	3.1	21.5	1.8	8.0
Trécé broad spectrum	---	---	108.6	---	108.0	12.9	8.1	3.4	14.7
Trécé selective	---	---	30.3	---	17.7	4.5	3.4	1.0	12.7

*=Numbers represent ratio of non-targets captured for each SWD captured; hence, a value of 1 would indicate one SWD captured for each non-target insect (= high selectivity). Conversely, high values indicate less selectivity for SWD. For each week, the most selective treatment (= the one with the lowest ratio) is highlighted in green.




Figure 1. Early-season (11 May to 8 July, 2020) captures of male and female spotted wing drosophila (SWD) in traps baited with diluted grape juice (DGJ) and four commercial lures, in four Massachusetts locations. Bars superscribed by the same letter are not significantly different at odds of 19:1.

represent greater number of SWD captured relative to non-targets) in 5 out of the 6 dates for which such ratios were calculated. On June 1, the only treatment that attracted SWD was DGJ, therefore no comparisons of ratios could be made. The Scentry lure had the best selectivity on the Jun 10 trapping date.

Figure 1 shows that, across the entire trapping period (May 11 to July 8) DGJ was a very attractive material to adult SWD. More specifically, DGJ captured 5.5, 2.2, 1.6, and 2.7 times more SWD than did Trécé -selective, Trécé -broad spectrum, Scentry, and AlphaScents lures, respectively (Figure 1). It is important to note that DGJ was replaced every week whereas the commercial lures were not replaced throughout the study. The expected lure longevity is 4-6 weeks for the two Trécé lures and for the Scentry lure, and 8 weeks for the AlphaScents lure.

Experiment 2: Effect of grape juice fermentation on SWD attraction. Aging DGJ for 1 or 2 weeks did no influence captures of male SWD in traps. In contrast, twice as many females were captured when DGJ was fermented for 1 or 2 weeks compared to fresh DGJ (Figure 2A). No statistical difference in captures was noted between the two levels of fermentation. Captures of non-target insects followed the same pattern




Figure 2. Captures (average number per trap) of (A) adult SWD (males and females combined) and (B) non-target insects (other insects belonging to the SWD family), according to treatment. This study was conducted at the UMass Cold Spring Orchard in 2019. For each SWD gender (top chart) and for non-targets (bottom chart), bars superscribed by the same letter are not significantly different at odds of 19:1.

described for female SWD. For instance, the number of other drosophilids was 70% and 74% greater when DGJ was fermented for 1 or 2 weeks, respectively, when compared to fresh DGJ (Figure 2B).

Conclusions

The results from the first study confirmed that DGJ is attractive to male and female SWD, it has high selectivity, and therefore this low-cost material can be used for monitoring SWD populations. Our fermentation study revealed that female captures can be increased if traps are left for up two weeks in the field (we did not evaluate longer intervals). However, the fermentation process will also attract more non-target insects. Further research will be conducted to determine ways in which fermentation can be manipulated so as to attract fewer non-target insects without affecting SWD captures.

Acknowledgments

We thank Tim Nourse, and Tom and Ben Clark for allowing us to work on their orchards. We also thank Emily Begonis for assistance. Funding for this research was provided by the Massachusetts Department of Agricultural Resources Specialty Crops Block Grant with funds provided by the United States Department of Agriculture (USDA), Agricultural Marketing Service through agreement number AM180100XXXXG025. The views or findings presented here are the Grantee's and do not necessarily represent those of the State or the Missouri Department of Agriculture.

References

Piñero, J.C. Foley, N. 2018. Evaluation of diluted grape juice as an inexpensive attractant for the invasive fruit pest spotted wing *Drosophila*. *Fruit Notes* 83(3) 1-7.

HAVE A BLAST WITH OUR SPRAYERS

REARS PUL-BLAST SPRAYERS
100-500 GALLON STAINLESS STEEL TANKS
Multiple profile, medium power, adaptive PTO machine. CV or standard hitch. Adaptable to many tractor sizes and power requirements.

Supplying Growers, Gardeners
OESCO, INC.
and Groundskeeping Professionals. Since 1954.

above: 200 gallon narrow Pul-Blast
below: 100 gallon Pul-Blast

300 gal. Pul-Blast Blueberry Special

Call for a catalog or just stop by **800-634-5557** www.oescoinc.com
8 Ashfield Rd./Rt. 116, P.O. Box 540, Conway, MA 01341

**THE LEADING
SMALL FRUIT NURSERY TODAY!**

NOURSE

The Best Berry Plants Since 1932

- *Excellent Customer Service*
- *Wide Variety Selection*
- *Technical Support*
- *Complete Lab Facility
for Tissue Culture
& Virus Indexing*



Strawberries Currants
Brambles Gooseberries
Asparagus Elderberries
Blueberries Rhubarb

NOURSE FARMS, INC
41 RIVER ROAD
SOUTH DEERFIELD MA 01373

413.665.2658

INFO@NOURSEFARMS.COM

NOURSEFARMS.COM

Eco-Friendly Insect, Disease, Bird Control

University/USDA tested

Stink Bug Traps

Brown Marmorated and Native Bugs

Insect Traps and Lures

*Plum Curculio Trap Tree **Control**,
Codling & Oriental Moth, Cranberry
Pests, **Black Stem Borer**, Others*

Honey Bee Lure

Attract Bees - Increase Pollination

Predalure attracts beneficials

Oriental Beetle MD

*Mating Disruption
Fruit Crops & Ornamentals*

Prestop

*New Biofungicide Impressive
Activity. Foliar/Root Diseases*

Avex

*Bird Control. Apply by ground or
air. Cherries, Blueberries, Sweet
Corn, other crops*



*Committed to the Environment and Green Technology
Since 1990*

P. 303-469-9221
agbio@agbio-inc.com
www.AgBio-Inc.com

