

# Accede™ Experimental Use Permit in Massachusetts in 2021 (and what to expect in 2022)

*Jon Clements, University of Massachusetts Extension*

*Duane Greene, Stockbridge School of Agriculture, University of Massachusetts*

In late 2020 Valent Biosciences received Federal registration for Accede™ plant growth regulator. From their [announcement](#) “Accede is the first PGR based on a naturally occurring compound developed specifically for thinning of stone fruit, including peaches and nectarines. It also gives apple growers an effective tool to thin apples in the late thinning window when fruit are 15-20 mm in diameter. Until now, no fruit thinner has provided reliable thinning at this stage of development. Use of Accede has the potential to reduce the need for costly hand thinning to adjust the crop load and generate higher fruit quality and grower returns.”

How does Accede work to thin apples and peaches? First, Accede is classified as a naturally occurring “biochemical” plant growth regulator. It is an immediate precursor to ethylene production. Application of Accede thus stimulates ethylene production which in turn triggers seed abortion and the formation of an abscission zone. Hence fruitlet drop (in apples when applied at 15-20 mm fruitlet diameter) and flower senescence (in peaches when applied at bloom) is promoted and accelerated. Application of ethephon can do the same thing, but ethephon is somewhat “unpredictable” whereas Accede appears to be “safer.”

At the University of Massachusetts, Duane Greene started experimenting with ACC (the active ingredient of Accede) over 10 years ago. A couple years of trialing at the UMass Orchard in Belchertown on McIntosh apples resulted in significant fruit thinning that increased (less percent fruit set) with rate and with later timing (10 mm vs. 20 mm, Figure 1). He also noted some leaf yellowing/drop after Accede application, but it was not excessive. Phil Schwallier at Michigan State University in 2016 showed that ACC application increased fruit

Effect of time and concentration of ACC application on thinning of McIntosh apples.		
Treatment (ppm)	Fruit/ cm LCSA	Percent set
1. Control 10 mm	10.2	83
2. ACC 200 10 mm	10.3	87
3. ACC 400 10 mm	4.6	39
4. Control 20 mm	10.7	87
5. ACC 200 20 mm	7.4	65
6. ACC 400 20 mm	1.4	13

**Figure 1.** 2011 results of ACC research at the UMass Cold Spring Orchard (Duane Greene).

size of Gala apples to 160 grams per apple compared to 120 grams per apple for untreated trees.

Lacking a state label in Massachusetts in 2021, we were asked by Valent to supervise Experimental Use Permit (EUP) applications of Accede in a few Massachusetts apple orchards. Valent applied for and received the EUP from the Massachusetts Department of Agricultural Resources (MDAR) in January 2021. Notable stipulations of the EUP included:

- MDAR notified in writing prior to application
- Supervised by Cat 49 (Research & Demonstration) applicator
- Public access limited by signage “Notice Pesticide Testing”
- Report submitted to MDAR at conclusion of EUP

With enough Accede on hand in the spring of 2021 to cover about five acres, I solicited four Massachusetts orchards – one in the northeast, two in central Massachusetts, and one west of the Connecticut River – to use Accede under the EUP. I also

intended to use Accede at the UMass Orchard in Belchertown. Along with Jim Wargo, Valent sales rep for New England and New York, visits were made to the orchards in late May to assess the need for further thinning after they had already applied their petal fall and 10 mm chemical thinners (Figure 2). It was not too difficult to identify one-half to one acre apple blocks where additional thinning was desired at 15-20 mm fruitlet size.



**Figure 2.** Accede EUP application discussion with Jim Wargo (Valent USA) on 25 May 2021.

Valent supplied the directions for the growers to use when applying Accede under the EUP, including:

- Use a rate of 200-400 ppm (23-46 fl. oz./A) at 15-20 mm fruitlet diameter
- Use a non-ionic surfactant at 0.05% v/v (6.5 fl oz/100 gal)
- Use 100 gallons per acre (adequate coverage of fruit and foliage)
- DO NOT apply as tank mix partner with other thinning products
- Consider reduced rate if temperatures predicted to exceed 90 F. on day of application
- And allow 7-10 days to observe effect of thinning

Growers were instructed to follow these directions the best they could, and applications were made around 1 June. We should note that the NEWA Apple Carbohydrate model indicated a moderate carbohydrate deficit would occur shortly after most applications were made, so we expected to get some (good) results. Accede was applied to several varieties, including Gala, Honeycrisp, Fuji, Mc- Intosh, Cortland, Macoun, Golden Delicious,

and Paulared (among a few others). Growers noted there were no problems mixing and applying Accede and most used LI-700 as the surfactant. The rate of Accede used was 34.5 to 46 fl. oz. in 50 to 100 gallons of water per acre. Except for Paulared and one Macoun block being larger trees, the rest of the blocks were smaller (150 gallons per acre dilute tree row volume approximately) but mature bearing trees on dwarfing rootstocks

A few weeks following the Accede applications, a visit was made to each orchard to visually assess the efficacy (or not) of the Accede application(s). Valent provided a form that included the assessment parameters of leaf yellowing/drop, tree vigor, thinning activity, and return bloom (Figure 3).

5. Assessments:

a. Leaf yellowing / Leaf drop: at 7 days after application, rate leaf yellowing or leaf drop on the following scale:

0 = None  
 1 = Slight  
 2 = Moderate  
 3 = Severe (commercially unacceptable)

b. Tree vigor: at 2-3 weeks after application, rate tree vigor or canopy density:

0 = Excellent (no difference in canopy density vs non-treated)  
 1 = Good (slight reduction in canopy density, but acceptable)  
 2 = Fair (noticeable reduction in canopy density)  
 3 = Poor (significant and unacceptable reduction in canopy density)

c. Thinning activity: at 3 to 4 weeks after application, but prior to hand thinning, visually rate thinning activity of ACCEDE:

0 = Under thinned (excessive hand thinning required)  
 1 = Sufficiently thinned (minimal to moderate hand thinning required)  
 2 = Over thinned

d. 2022 Return bloom: ACCEDE treated area vs non-treated area

0 = ACCEDE area has lower return bloom compared to non-treated  
 1 = ACCEDE area with same return bloom  
 2 = ACCEDE area with greater return bloom

**Figure 3.** Portion of assessment form provided by Valent to evaluate Accede EUP applications.

When assessments were complete, it was agreed that leaf yellowing/leaf drop ranged from none to moderate. Leaf yellowing typically shows up three to four days after application on older and smaller leaves in the spur. Yellowed leaves fall off rapidly so there is little evidence of the leaf yellowing on the tree after 10 days. Tree vigor was generally good (slight reduction in canopy density) to excellent (no difference in tree canopy vs. untreated). Thinning activity varied widely by orchard and variety. As a rule, Golden Delicious types (including Gala) were sufficiently thinned compared to the untreated control. (Figures 4 and 5), thus little or no hand thinning was needed. McIntosh types seemed largely unfazed



**Figure 4.** Gala fruitlet drop following Accede application: untreated control on left vs. Accede application on right.



**Figure 5.** Typical Gala fruitlet clusters after Accede application: untreated control on left, Accede application on right.

by the Accede application(s). This was also observed in an ACC experiment conducted on McIntosh by Greene in 2021 where virtually no thinning occurred too. On Honeycrisp, it's unclear how Accede may work, it may be useful but this remains to be determined.

We are learning how Accede may be another useful chemical thinner in your toolbox but don't count on it being a silver bullet. Plan on starting chemical thinning early and often just as usual, and then use Accede if you get to the point when fruitlets are 15-20 mm in size and it appears that you will have to do considerable hand thinning. Accede will probably not be a good option at the 10 mm fruit size, but that is still being evaluated. Having some carbohydrate deficit on the trees during the timing and 3-4 days following Accede application will be helpful (and probably necessary). Accede may be particularly useful on Gala-type apples to increase thinning and improve apple size.

We alluded to the fact that Accede can also be used for thinning stone fruit, including peaches! That is a big deal as we don't have a chemical thinner for peaches. And it's a very promising thinner of peaches. For example, a summary of Accede EUP applied


to peaches in 2021 in New York, Michigan, Ohio, and Pennsylvania across many peach varieties resulted in an average fruit reduction of 39% compared to the untreated control (Gregory Clarke, Valent USA). Peach varieties did vary quite widely in their thinning response to Accede, but that might have been a location effect. In Michigan, work by Anna Wallis and Phil Schwallier in 2021 showed that hand thinning time was reduced by approximately 50% with Accede application compared to non-treated peaches. That would be sweet!

New York already has a state registration for Accede, on both apples and stone fruit, but as of late 2021 not all the New England states had Accede registered, including Massachusetts (per [CDMS Label Search](#)). All state registrations, however, are expected in 2022. We have learned that Accede supply for 2022 may be somewhat limited, and Valent plans to expand a demo program like what was done in 2021. Reach out to your agrochemical supplier if you want to use Accede in 2022, particularly if you want to trial it on some peaches.

### *Acknowledgments*

Thanks to Parlee Farms, Carlson Orchards, Ragged Hill Orchard, and Clark Brothers Orchard for participating in the EUP. And to Jim Wargo and Greg Clarke of Valent for supporting the EUP in more ways than one.





# Fresher, Sweeter, Juicier

*Peaches From Circle M Farms*



*Exceptional quality and flavor at attractive prices  
Brought to you from the 4th generation family  
farm of Santo John Maccherone*



- PLU Stickers Available Upon Request
- Low in Calories
- Nutritious and Delicious
- Exceptional Flavor



## SANTO SAYS...

"Peaches from Circle M Farms just plain taste better. Their size and color are what your customers want, and their flavor is the icing on the cake.

That's because our orchards are tended with care and attention, and their fruit is left on the tree to ripen naturally until they reach their optimum sweetness. Then they are picked by hand, taste-tested by me to ensure quality, and packed gently into

25 lb. volume boxes for shipping. Single layer cartons are available upon request.

Circle M's 30 varieties of peaches and nectarines are known for their sweetness and their luscious color. They look as good as they taste.

Discriminating customers appreciate our white flesh peaches because of their lower acid levels and supremely sweet taste.

Add it all up: Peaches from Circle M Farms are fresher and sweeter, picked by hand when their time has come, and packed with care for a longer shelf life.

Yellow, white and donut peaches and yellow and white nectarines are available early July to late September."



Call us at 856 856 207-4542 or  
609 381-6465. Fax your order to  
856 878-0404



Available in 16 oz. and 64 oz. sizes

**Made from fresh New Jersey Peaches  
"Peach Cider Drink, Peach Salsa, Peach  
Preserves"**

Santo John Maccherone  
[circlemfarmsllc@gmail.com](mailto:circlemfarmsllc@gmail.com)

**Circle M Farms  
88 Sandy Ridge Road  
Salem, N.J. 08079**



Adams County Nursery, Inc.

**Growers of Fruit Tree Nursery Stock**

26 Nursery Road, Aspers, PA 17304

acn@acnursery.com • 717.677.8105 • www.acnursery.com

